



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

**Coffein and theobromin.**—In a detailed account of his studies on the genera *Coffea*, *Thea*, *Theobroma*, and *Cola*, WEEVERS answers for the first time these questions:<sup>25</sup> Do coffein and theobromin arise in connection with the proteid metabolism of the plant or not? If so, do they arise through analysis or synthesis of proteids, and can they be used in proteid synthesis? He concludes that both are produced in consequence of secondary processes of dissimilation, that they are stored for a longer or shorter time, and then are again used in proteid synthesis. They constitute a very compact means of storing N, as they contain about 30 per cent. N as compared with 19 per cent. or less in proteids, and this is doubtless the reason why they occur so abundantly in seeds, though they may have also a protective function.—C. R. B.

**Desert vegetation in South Africa.**—The country of Welwitschia has always seemed fascinating to botanical travelers. Its accessibility is not fully realized; not only is it near the coast, but the heart of the desert is traversed by a railroad which has a station named Welwitsch from the abundance of the remarkable gymnosperm in the region thereabouts. PEARSON<sup>26</sup> has given an account of a journey through a part of German Southwest Africa, in which there is a description of the coastal strip with *Ancathosicyos horrida*, the Namib plains with Welwitschia, and the dry river beds with a less xerophytic vegetation closely resembling the Acacia vegetation farther east. Near Windhuk, the seat of government, there are Acacia park forests, in which *Acacia giraffae* is the predominating species.—H. C. COWLES.

**Root-parasitism.**—BARBER<sup>27</sup> has continued his studies of the haustorium of *Santalum album*. In the first paper<sup>28</sup> the early stages up to penetration were described; while the present paper deals with the mature haustorium. The penetration of the haustorium, its general structure, and its relation to the host tissues are first described. Then follow detailed descriptions of the cortex and "nucleus," the vascular system and its connections with the host, the character and development of the "sucker" with its double function of secretion and absorption, cases of irregular penetration, the general activity of the haustorium, and the interrelations of host and parasite. The numerous plates serve to make the descriptions remarkably clear.—J. M. C.

**Taiwania.**—HAYATA described this new genus of Coniferales from the Island of Formosa in 1906,<sup>29</sup> from a few dry branches bearing cones. Since that time he

<sup>25</sup> WEEVERS, TH., Die physiologische Bedeutung des Koffeins und des Theobromins. Ann. Jard. Bot. Buitenzorg 21:1-78 1907.

<sup>26</sup> PEARSON, H. H. W., Some notes on a journey from Walfish Bay to Windhuk. Kew Bull. 1907:339-360.

<sup>27</sup> BARBER, C. A., Studies in root-parasitism. The haustorium of *Santalum album*. II. Mem. Depart. Agric. India Bot. 1:no. 12, pp. 58. pls. 15. 1907.

<sup>28</sup> BOT. GAZETTE 40:159. 1905 and 42:317. 1906.

<sup>29</sup> Jour. Linn. Soc. Bot. 32:330-332. pl. 16.